

ADULT BEDSIDE MODERATE SEDATION PLAN
- Phase: Intra-Procedure Orders

PHYSICIAN ORDERS

Diagnosis _____

Weight _____ Allergies _____

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER ORDER DETAILS

Patient Care

Vital Signs

T;N, Per Policy

Patient Position

Lying on Left Side

Prone

Other

Lying on Right Side

Supine

Oxygen Administration

Via: Nasal cannula, Keep sats greater than: 90%

Medications

Medication sentences are per dose. You will need to calculate a total daily dose if needed.

Other Medications

lidocaine (lidocaine 1% injectable solution)

1 mL, locally, inj, ONE TIME

Reversal Agent

flumazenil

0.2 mg, IVPush, inj, q1min, PRN excess sedation

naloxone

0.2 mg, IVPush, inj, q2min, PRN bradypnea

TO Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: _____ Date _____ Time _____

Physician Signature: _____ Date _____ Time _____



UMC Health System

Patient Label Here

ADULT BEDSIDE MODERATE SEDATION PLAN
- Phase: Medication Documentation

PHYSICIAN ORDERS

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER ORDER DETAILS

...Additional Orders

TO Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: _____ Date _____ Time _____

Physician Signature: _____ Date _____ Time _____



ADULT BEDSIDE MODERATE SEDATION PLAN
- Phase: Post-Procedure Orders

PHYSICIAN ORDERS

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER	ORDER DETAILS
Patient Care	
Patient Activity <input type="checkbox"/> Bedrest <input type="checkbox"/> Bedrest, Bed Position: HOB Flat <input type="checkbox"/> Up Ad Lib/Activity as Tolerated	
Communication	
Notify Nurse (DO NOT USE FOR MEDS)	
Confirm Line Placement - Cleared for Use (Cleared for Use - CVL)	
Diagnostic Tests	
DX Chest Single View	

TO Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: _____ Date _____ Time _____

Physician Signature: _____ Date _____ Time _____



ADULT BEDSIDE MODERATE SEDATION PLAN
- Phase: Pre-Procedure Orders

PHYSICIAN ORDERS

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER	ORDER DETAILS
	Patient Care
	Sedation History and Physical Update
	Moderate Sedation Medications at Bedside
	Set Up for Biopsy
	Set Up for Bronchoscopy
	Set Up for Cardioversion - Chemical
	Set Up for Cast Application
	Set Up for Central Line Placement
	Set Up for Chest Tube
	Set Up for Closed Reduction
	Set Up for Fracture Manipulation
	Set Up for Incision and Drainage at Beds (Set Up for Incision and Drainage at Bedside)
	Set Up for Laceration Repair
	Set Up for Lumbar Puncture
	Set Up for Paracentesis
	Set Up For Pleural Catheter
	Set Up for Thoracentesis
	Set Up for Tracheostomy
	Communication
	Notify Nurse (DO NOT USE FOR MEDS) <input type="checkbox"/> T;N, Ensure IV is patent
	Obtain Consent <input type="checkbox"/> T;N, Consent for: Moderate sedation

TO Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: _____ Date _____ Time _____

Physician Signature: _____ Date _____ Time _____



ADULT BEDSIDE MODERATE SEDATION PLAN
- Phase: NURSING SEDATION MEDICATION
DOCUMENTATION PLAN

PHYSICIAN ORDERS

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER	ORDER DETAILS
Communication	
This plan is for nursing documentation only. This plan should only be entered by nurses for medication documentation following a procedure. Procedural Sedation Medications Guidelin (Procedural Sedation Medications Guidelines)	
Medications	
Medication sentences are per dose. You will need to calculate a total daily dose if needed.	
Analgesics: Opioids	
	fentaNYL <input type="checkbox"/> 25 mcg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication. <input type="checkbox"/> 50 mcg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication. <input type="checkbox"/> 75 mcg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication. <input type="checkbox"/> 100 mcg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication.
	HYDRomorphone <input type="checkbox"/> 0.5 mg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication. <input type="checkbox"/> 1 mg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication. <input type="checkbox"/> 2 mg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication.
	morphine <input type="checkbox"/> 2 mg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication. <input type="checkbox"/> 4 mg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication. <input type="checkbox"/> 10 mg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication.
Sedatives	
	LORazepam <input type="checkbox"/> 0.5 mg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication. <input type="checkbox"/> 1 mg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication. <input type="checkbox"/> 2 mg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication.
	midazolam <input type="checkbox"/> 0.5 mg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication. <input type="checkbox"/> 2 mg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication. <input type="checkbox"/> 4 mg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication. <input type="checkbox"/> 5 mg, IVPush, inj, OCTOR For Procedural Sedation Only. See INet for incremental dosing of this medication.

TO Read Back Scanned Powerchart Scanned PharmScan

Order Taken by Signature: _____ Date _____ Time _____

Physician Signature: _____ Date _____ Time _____



ADULT BEDSIDE MODERATE SEDATION PLAN
- Phase: NURSING SEDATION MEDICATION
DOCUMENTATION PLAN

PHYSICIAN ORDERS

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER	ORDER DETAILS
	<p>diphenhydrAMINE <input type="checkbox"/> 25 mg, IVPush, inj, OCTOR</p>
	<p>The Following Medications are for Deep Sedation Use Only</p> <p>dexmedeTOMIDine (dexmedetomidine 20 mcg/10 mL) <input type="checkbox"/> 25 mcg, Slow IVPush, inj, OCTOR For Deep Sedation Use Only. See INet for incremental dosing of this medication during procedural sedation. <input type="checkbox"/> 50 mcg, Slow IVPush, inj, OCTOR For Deep Sedation Use Only. See INet for incremental dosing of this medication during procedural sedation. <input type="checkbox"/> 100 mcg, Slow IVPush, inj, OCTOR For Deep Sedation Use Only. See INet for incremental dosing of this medication during procedural sedation.</p>
	<p>etomidate <input type="checkbox"/> 5 mg, IVPush, inj, OCTOR For Deep Sedation Use Only. See INet for incremental dosing of this medication during procedural sedation. <input type="checkbox"/> 10 mg, IVPush, inj, OCTOR For Deep Sedation Use Only. See INet for incremental dosing of this medication during procedural sedation. <input type="checkbox"/> 20 mg, IVPush, inj, OCTOR For Deep Sedation Use Only. See INet for incremental dosing of this medication during procedural sedation.</p>
	<p>ketamine <input type="checkbox"/> 50 mg, IVPush, inj, OCTOR For Deep Sedation Use Only. See INet for incremental dosing of this medication during procedural sedation. <input type="checkbox"/> 100 mg, IVPush, inj, OCTOR For Deep Sedation Use Only. See INet for incremental dosing of this medication during procedural sedation. <input type="checkbox"/> 200 mg, IVPush, inj, OCTOR For Deep Sedation Use Only. See INet for incremental dosing of this medication during procedural sedation.</p>
	<p>propofol <input type="checkbox"/> 10 mg, IVPush, inj, OCTOR For Deep Sedation Use Only. See INet for incremental dosing of this medication during procedural sedation. <input type="checkbox"/> 20 mg, IVPush, inj, OCTOR For Deep Sedation Use Only. See INet for incremental dosing of this medication during procedural sedation. <input type="checkbox"/> 50 mg, IVPush, inj, OCTOR For Deep Sedation Use Only. See INet for incremental dosing of this medication during procedural sedation. <input type="checkbox"/> 100 mg, IVPush, inj, OCTOR For Deep Sedation Use Only. See INet for incremental dosing of this medication during procedural sedation.</p>

TO Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: _____ Date _____ Time _____

Physician Signature: _____ Date _____ Time _____

