

<p>UMC Health System</p> <p>ADULT BOWEL PROGRAM PLAN</p>	<p>Patient Label Here</p>
--	----------------------------------

PHYSICIAN ORDERS

Diagnosis _____

Weight	_____	Allergies	_____
---------------	-------	------------------	-------

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER ORDER DETAILS

Communication

This plan is to rule out intestinal obstruction.

Medications

Medication sentences are per dose. You will need to calculate a total daily dose if needed.

Start patient on Option #1. If no results within twelve hours, continue Option #1 and implement Option #2. If no results within six hours, continue Option #1 and implement Option #3. If no results after six hours or if continuing diarrhea or severe cramping is experienced, contact physician.

Bowel Preparation: Option #1

docusate	<input type="checkbox"/> 100 mg, PO, cap, BID	Do not crush or chew.
	<input type="checkbox"/> 100 mg, PO, liq, BID	

bisacodyl	<input type="checkbox"/> 10 mg, PO, tab ec, Nightly	Do not crush or chew.
------------------	---	-----------------------

senna	<input type="checkbox"/> 17.2 mg, PO, tab, Nightly	<input type="checkbox"/> 17.6 mg, per tube, liq, Nightly, [17.6 mg/10 mL]
--------------	--	---

Bowel Preparation: Option #2

bisacodyl	<input type="checkbox"/> 10 mg, rectally, supp, ONE TIME	Give rectal bisacodyl if no results from docusate, bisacodyl oral, or senna after 12 hours.
------------------	--	---

Bowel Preparation: Option #3

sodium biphosphate-sodium phosphate (Fleet Enema)	<input type="checkbox"/> 1 ea, rectally, enema, ONE TIME	Give Fleet enema if no results from rectal bisacodyl after six hours.
--	--	---

--	--	--

TO Read Back Scanned Powerchart Scanned PharmScan

Order Taken by Signature: _____ Date _____ Time _____

Physician Signature: _____ Date _____ Time _____

