

<b>UMC Health System</b>  ALTEPLASE INTRA-PLEURAL PLAN	Patient Label Here
--	--------------------

**PHYSICIAN ORDERS**

**Diagnosis** \_\_\_\_\_

**Weight** \_\_\_\_\_ **Allergies** \_\_\_\_\_

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

**ORDER ORDER DETAILS**

**Medications**

**Medication sentences are per dose. You will need to calculate a total daily dose if needed.**

Select medication orders below based on syringe type needed:

**CATHETER TIP SYRINGE**

**alteplase (alteplase intra-pleural)**

10 mg, intra-pleural, q12h, x 6 dose, \*\*CATHETER TIP SYRINGE, intra-pleural use ONLY\*\*  
 \*\*Prepare in 60 mL catheter tip syringe\*\* Allow to dwell for 1 hour, then drain for 1 hour. If ordered, administer dornase 2 hours AFTER alteplase dose.

5 mg, intra-pleural, q12h, x 6 dose, \*\*CATHETER TIP SYRINGE, intra-pleural use ONLY\*\*  
 \*\*Prepare in 60 mL catheter tip syringe\*\* Allow to dwell for 1 hour, then drain for 1 hour. If ordered, administer dornase 2 hours AFTER alteplase dose.

10 mg, intra-pleural, q24h, x 3 dose, \*\*CATHETER TIP SYRINGE, intra-pleural use ONLY\*\*  
 \*\*Prepare in 60 mL catheter tip syringe\*\* Allow to dwell for 1 hour, then drain for 1 hour. If ordered, administer dornase 2 hours AFTER alteplase dose.

5 mg, intra-pleural, q24h, x 3 dose, \*\*CATHETER TIP SYRINGE, intra-pleural use ONLY\*\*  
 \*\*Prepare in 60 mL catheter tip syringe\*\* Allow to dwell for 1 hour, then drain for 1 hour. If ordered, administer dornase 2 hours AFTER alteplase dose.

**dornase alfa (dornase alfa intra-pleural)**

5 mg, intra-pleural, q12h, x 6 dose, \*\*CATHETER TIP SYRINGE, intra-pleural use ONLY\*\* If ordered, administer dornase 2 hours AFTER alteplase dose.  
 \*\*Prepare in 60 mL catheter tip syringe\*\* Administer 2 hours AFTER alteplase, IF ordered. Allow to dwell for 1 hour, then drain for 1 hour.

5 mg, intra-pleural, q24h, x 3 dose, \*\*CATHETER TIP SYRINGE, intra-pleural use ONLY\*\* If ordered, administer dornase 2 hours AFTER alteplase dose.  
 \*\*Prepare in 60 mL catheter tip syringe\*\* Administer 2 hours AFTER alteplase, IF ordered. Allow to dwell for 1 hour, then drain for 1 hour.

**LUER-LOCK SYRINGE**

**alteplase (alteplase intra-pleural)**

10 mg, intra-pleural, q12h, x 6 dose, \*\*LUER-LOCK SYRINGE, intra-pleural use ONLY\*\*  
 \*\*Prepare in 60 mL luer-lock syringe\*\* Send med request when ready for dose. Allow to dwell for 1 hour, then drain for 1 hour. If ordered, administer dornase 2 hours AFTER alteplase dose.

5 mg, intra-pleural, q12h, x 6 dose, \*\*LUER-LOCK SYRINGE, intra-pleural use ONLY\*\*  
 \*\*Prepare in 60 mL luer-lock syringe\*\* Send med request when ready for dose. Allow to dwell for 1 hour, then drain for 1 hour. If ordered, administer dornase 2 hours AFTER alteplase dose.

Continued on next page....

TO  Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Physician Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_



PHYSICIAN ORDERS

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER	ORDER DETAILS
	<input type="checkbox"/> 10 mg, intra-pleural, q24h, x 3 dose, **LUER-LOCK SYRINGE, intra-pleural use ONLY** **Prepare in 60 mL luer-lock syringe** Send med request when ready for dose. Allow to dwell for 1 hour, then drain for 1 hour. If ordered, administer dornase 2 hours AFTER alteplase dose. <input type="checkbox"/> 5 mg, intra-pleural, q24h, x 3 dose, **LUER-LOCK SYRINGE, intra-pleural use ONLY** **Prepare in 60 mL luer-lock syringe** Send med request when ready for dose. Allow to dwell for 1 hour, then drain for 1 hour. If ordered, administer dornase 2 hours AFTER alteplase dose.
	<p><b>dornase alfa (dornase alfa intra-pleural)</b></p> <input type="checkbox"/> 5 mg, intra-pleural, q12h, x 6 dose, **LUER-LOCK SYRINGE, intra-pleural use ONLY** If ordered, administer dornase 2 hours AFTER alteplase dose. **Prepare in 60 mL luer-lock syringe** Administer 2 hours AFTER alteplase, IF ordered. Allow to dwell for 1 hour, then drain for 1 hour. <input type="checkbox"/> 5 mg, intra-pleural, q24h, x 3 dose, **LUER-LOCK SYRINGE, intra-pleural use ONLY** If ordered, administer dornase 2 hours AFTER alteplase dose. **Prepare in 60 mL luer-lock syringe** Administer 2 hours AFTER alteplase, IF ordered. Allow to dwell for 1 hour, then drain for 1 hour.

TO  Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Physician Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

