



UMC HEALTH SYSTEM

ANESTHESIA PRE-OPERATIVE PLAN

Patient Label Here

A UMC Health System Performance Improvement Initiative for use in all units where patients for surgical procedures

*Denotes guideline requirement for Core Measures

- 1. ALLERGIES: NKDA Allergic to:
2. CONSULT:
3. NURSING: Weight in kilograms Old chart to OPS/OR Holding area for review by anesthesiologist Verify last oral intake status Pre-Operative warming to keep patient at a temp >= 97 F method every 15 minutes times 2 (over age 16 years)
4. LABORATORY/DIAGNOSTICS: Urine Pregnancy Test day of surgery (All childbearing bearing age females) Initiate Pre-Operative Testing Guidelines based on Expected Surgical Blood Loss, Patient age, Medical Condition, Medications Point of Care Glucose for Diabetic Patients: please call anesthesiologist for blood sugar > blood sugar < EKG Chest X-ray
5. IV: Adult: 1000 milliliters Lactated Ringers at 100 milliliters/hour on arrival to OPS (Renal Failure Patients check with MD) Pediatric: No intravenous access; Intravenous will be started by physician in the OR for age 10 and below Continuous intravenous fluids to run at milliliters/hour
6. MEDICATIONS: Refer also to Admission Medication Reconciliation Form Pain Ease spray to intended site for Intravenous. In holding give: Adult: Famotidine (Pepcid) 20 milligrams, intravenous on call to OR Reglan 10 milligrams, intravenous on call to OR Bicitra 30 milliliters, by mouth on call to OR (do not give for Ophthalmology Procedures) Acetaminophen /Hydrocodone (Lortab elixir) 10 - 15 ml PO on call in OR Holding Child: Versed (0.25 milligram - 0.5milligrams/kilogram) = by mouth on call to OR (maximum dose 20 milligrams) Acetaminophen/Hydrocodone (Lortab elixir) 0.2 -0.4 ml/kg/dose = ml PO on call in Holding (Children 2-13 or < 50 kg) Ibuprofen Suspension (Children's Motrin) 5 milligrams to 10 milligrams/kilogram by mouth on call to OR

TO Read back Order taken by Signature: Date/Time:

Physician Signature Date/Time

