

DIANEAL 2.5% WITH CEFTAZIDIME 1G NIGHTLY - VANCOMYCIN 1G EVERY 4TH NIGHT

PHYSICIAN ORDERS

Diagnosis \_\_\_\_\_

Weight \_\_\_\_\_

Allergies \_\_\_\_\_

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER ORDER DETAILS

IV Solutions

ALL of the components below are REQUIRED to complete the following regimen: Dianeal 2.5% four times daily with cefTAZidime 1g nightly AND vancomycin 1g every 4th night.

Dex 2.5% + lytes (Dianeal PD-2) (Dianeal PD-2 with 2.5% Dextrose intraperitoneal solution)

2,000 mL, intra-peritoneal, dialysate, TID, Infuse over 0 min, Dwell Volume: 2,000

dianeal 2.5% + vancomycin 1 g + cefTAZad (dianeal 2.5% + vancomycin 1 g + cefTAZadime 1 g)

1 g, intra-peritoneal, dialysate, q96h, Dwell Volume: 2,000 mLs

1 g, Every Bag

2,000 mL, Every Bag

dianeal 2.5% + cefTAZidime 1g

1 g, intra-peritoneal, dialysate, q96h, Dwell Volume: 2,000 mLs

2,000 mL, Every Bag

dianeal 2.5% + cefTAZidime 1g

2,000 mL, Every Bag

1 g, intra-peritoneal, dialysate, q96h, Dwell Volume: 2,000 mLs

dianeal 2.5% + cefTAZidime 1g

1 g, intra-peritoneal, dialysate, q96h, Dwell Volume: 2,000 mLs

2,000 mL, Every Bag

TO  Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Physician Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

