



<b>UMC Health System</b>  SLIDING SCALE INSULIN ASPART PLAN	Patient Label Here
---	--------------------

**PHYSICIAN ORDERS**

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER	ORDER DETAILS
-------	---------------

**Patient Care**

<b>POC Blood Sugar Check</b> <input type="checkbox"/> Per Sliding Scale Insulin Frequency <input type="checkbox"/> AC & HS 3 days <input type="checkbox"/> BID <input type="checkbox"/> q6h <input type="checkbox"/> q4h	<input type="checkbox"/> AC & HS <input type="checkbox"/> TID <input type="checkbox"/> q12h <input type="checkbox"/> q6h 24 hr
---	---

<b>Sliding Scale Insulin Aspart Guidelines</b> <input type="checkbox"/> Follow SSI Aspart Reference Text	
---	--

**Medications**

**Medication sentences are per dose. You will need to calculate a total daily dose if needed.**

<b>insulin aspart (Low Dose Insulin Aspart Sliding Scale)</b> <input type="checkbox"/> 0-10 units, subcut, inj, AC & nightly, PRN glucose levels - see parameters Low Dose Insulin Aspart Sliding Scale If blood glucose is less than 70 mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.  70-150 mg/dL - 0 units 151-200 mg/dL - 1 units subcut 201-250 mg/dL - 2 units subcut 251-300 mg/dL - 3 units subcut 301-350 mg/dL - 4 units subcut 351-400 mg/dL - 6 units subcut  If blood glucose is greater than 400 mg/dL administer 10 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. Once blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale. <input type="checkbox"/> 0-10 units, subcut, inj, BID, PRN glucose levels - see parameters Low Dose Insulin Aspart Sliding Scale If blood glucose is less than 70 mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.  70-150 mg/dL - 0 units 151-200 mg/dL - 1 units subcut 201-250 mg/dL - 2 units subcut 251-300 mg/dL - 3 units subcut 301-350 mg/dL - 4 units subcut 351-400 mg/dL - 6 units subcut  If blood glucose is greater than 400 mg/dL administer 10 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. Once blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale. Continued on next page....	
--	--

<input type="checkbox"/> TO <input type="checkbox"/> Read Back	<input type="checkbox"/> Scanned Powerchart	<input type="checkbox"/> Scanned PharmScan	
Order Taken by Signature: _____	Date: _____	Time: _____	
Physician Signature: _____	Date: _____	Time: _____	



SLIDING SCALE INSULIN ASPART PLAN

PHYSICIAN ORDERS

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER	ORDER DETAILS
	<p><input type="checkbox"/> 0-10 units, subcut, inj, TID, PRN glucose levels - see parameters                      Low Dose Insulin Aspart Sliding Scale                      If blood glucose is less than 70 mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.</p> <p>70-150 mg/dL - 0 units                      151-200 mg/dL - 1 units subcut                      201-250 mg/dL - 2 units subcut                      251-300 mg/dL - 3 units subcut                      301-350 mg/dL - 4 units subcut                      351-400 mg/dL - 6 units subcut</p> <p>If blood glucose is greater than 400 mg/dL administer 10 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. Once blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale.</p> <p><input type="checkbox"/> 0-10 units, subcut, inj, q6h, PRN glucose levels - see parameters                      Low Dose Insulin Aspart Sliding Scale                      If blood glucose is less than 70 mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.</p> <p>70-150 mg/dL - 0 units                      151-200 mg/dL - 1 units subcut                      201-250 mg/dL - 2 units subcut                      251-300 mg/dL - 3 units subcut                      301-350 mg/dL - 4 units subcut                      351-400 mg/dL - 6 units subcut</p> <p>If blood glucose is greater than 400 mg/dL administer 10 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. Once blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale.</p> <p><input type="checkbox"/> 0-10 units, subcut, inj, q4h, PRN glucose levels - see parameters                      Low Dose Insulin Aspart Sliding Scale                      If blood glucose is less than 70 mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.</p> <p>70-150 mg/dL - 0 units                      151-200 mg/dL - 1 units subcut                      201-250 mg/dL - 2 units subcut                      251-300 mg/dL - 3 units subcut                      301-350 mg/dL - 4 units subcut                      351-400 mg/dL - 6 units subcut</p> <p>If blood glucose is greater than 400 mg/dL administer 10 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. Once blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale.</p> <p>Continued on next page....</p>

TO  Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Physician Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_



SLIDING SCALE INSULIN ASPART PLAN

PHYSICIAN ORDERS

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER	ORDER DETAILS
	<p><b>insulin aspart (Moderate Dose Insulin Aspart Sliding Scale)</b></p> <p><input type="checkbox"/> 0-12 units, subcut, inj, AC &amp; nightly, PRN glucose levels - see parameters                      Moderate Dose Insulin Aspart Sliding Scale                      If blood glucose is less than 70mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.</p> <p>70-150 mg/dL - 0 units                      151-200 mg/dL - 2 units subcut                      201-250 mg/dL - 3 units subcut                      251-300 mg/dL - 5 units subcut                      301-350 mg/dL - 7 units subcut                      351-400 mg/dL - 10 units subcut</p> <p>If blood glucose is greater than 400 mg/dL, administer 12 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. Once blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale.</p> <p><input type="checkbox"/> 0-12 units, subcut, inj, BID, PRN glucose levels - see parameters                      Moderate Dose Insulin Aspart Sliding Scale                      If blood glucose is less than 70mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.</p> <p>70-150 mg/dL - 0 units                      151-200 mg/dL - 2 units subcut                      201-250 mg/dL - 3 units subcut                      251-300 mg/dL - 5 units subcut                      301-350 mg/dL - 7 units subcut                      351-400 mg/dL - 10 units subcut</p> <p>If blood glucose is greater than 400 mg/dL, administer 12 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. Once blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale.</p> <p><input type="checkbox"/> 0-12 units, subcut, inj, TID, PRN glucose levels - see parameters                      Moderate Dose Insulin Aspart Sliding Scale                      If blood glucose is less than 70mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.</p> <p>70-150 mg/dL - 0 units                      151-200 mg/dL - 2 units subcut                      201-250 mg/dL - 3 units subcut                      251-300 mg/dL - 5 units subcut                      301-350 mg/dL - 7 units subcut                      351-400 mg/dL - 10 units subcut</p> <p>If blood glucose is greater than 400 mg/dL, administer 12 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. Once blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale.</p> <p>Continued on next page....</p>

TO  Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Physician Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_



SLIDING SCALE INSULIN ASPART PLAN

PHYSICIAN ORDERS

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER	ORDER DETAILS
	<p><input type="checkbox"/> 0-12 units, subcut, inj, q6h, PRN glucose levels - see parameters                      Moderate Dose Insulin Aspart Sliding Scale                      If blood glucose is less than 70mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.</p> <p>70-150 mg/dL - 0 units                      151-200 mg/dL - 2 units subcut                      201-250 mg/dL - 3 units subcut                      251-300 mg/dL - 5 units subcut                      301-350 mg/dL - 7 units subcut                      351-400 mg/dL - 10 units subcut</p> <p>If blood glucose is greater than 400 mg/dL, administer 12 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. Once blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale.</p> <p><input type="checkbox"/> 0-12 units, subcut, inj, q4h, PRN glucose levels - see parameters                      Moderate Dose Insulin Aspart Sliding Scale                      If blood glucose is less than 70mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.</p> <p>70-150 mg/dL - 0 units                      151-200 mg/dL - 2 units subcut                      201-250 mg/dL - 3 units subcut                      251-300 mg/dL - 5 units subcut                      301-350 mg/dL - 7 units subcut                      351-400 mg/dL - 10 units subcut</p> <p>If blood glucose is greater than 400 mg/dL, administer 12 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. Once blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale.</p>
	<p><b>insulin aspart (High Dose Insulin Aspart Sliding Scale)</b></p> <p><input type="checkbox"/> 0-14 units, subcut, inj, AC &amp; nightly, PRN glucose levels - see parameters                      High Dose Insulin Aspart Sliding Scale                      If blood glucose is less than 70 mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.</p> <p>70-150 mg/dL - 0 units                      151-200 mg/dL - 3 units subcut                      201-250 mg/dL - 5 units subcut                      251-300 mg/dL - 7 units subcut                      301-350 mg/dL - 10 units subcut                      351-400 mg/dL - 12 units subcut</p> <p>If blood glucose is greater than 400mg/dL, administer 14 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. Once blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale.</p> <p>Continued on next page....</p>

TO  Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Physician Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_



SLIDING SCALE INSULIN ASPART PLAN

PHYSICIAN ORDERS

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER	ORDER DETAILS
	<p><input type="checkbox"/> 0-14 units, subcut, inj, BID, PRN glucose levels - see parameters  High Dose Insulin Aspart Sliding Scale  If blood glucose is less than 70 mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.</p> <p>70-150 mg/dL - 0 units  151-200 mg/dL - 3 units subcut  201-250 mg/dL - 5 units subcut  251-300 mg/dL - 7 units subcut  301-350 mg/dL - 10 units subcut  351-400 mg/dL - 12 units subcut</p> <p>If blood glucose is greater than 400mg/dL, administer 14 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. ONce blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale.</p> <p><input type="checkbox"/> 0-14 units, subcut, inj, TID, PRN glucose levels - see parameters  High Dose Insulin Aspart Sliding Scale  If blood glucose is less than 70 mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.</p> <p>70-150 mg/dL - 0 units  151-200 mg/dL - 3 units subcut  201-250 mg/dL - 5 units subcut  251-300 mg/dL - 7 units subcut  301-350 mg/dL - 10 units subcut  351-400 mg/dL - 12 units subcut</p> <p>If blood glucose is greater than 400mg/dL, administer 14 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. ONce blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale.</p> <p><input type="checkbox"/> 0-14 units, subcut, inj, q6h, PRN glucose levels - see parameters  High Dose Insulin Aspart Sliding Scale  If blood glucose is less than 70 mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.</p> <p>70-150 mg/dL - 0 units  151-200 mg/dL - 3 units subcut  201-250 mg/dL - 5 units subcut  251-300 mg/dL - 7 units subcut  301-350 mg/dL - 10 units subcut  351-400 mg/dL - 12 units subcut</p> <p>If blood glucose is greater than 400mg/dL, administer 14 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. ONce blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale.</p> <p>Continued on next page....</p>

TO  Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Physician Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_



SLIDING SCALE INSULIN ASPART PLAN

PHYSICIAN ORDERS

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER	ORDER DETAILS
	<input type="checkbox"/> 0-14 units, subcut, inj, q4h, PRN glucose levels - see parameters High Dose Insulin Aspart Sliding Scale If blood glucose is less than 70 mg/dL and patient is symptomatic, initiate hypoglycemia guidelines and notify provider.  70-150 mg/dL - 0 units 151-200 mg/dL - 3 units subcut 201-250 mg/dL - 5 units subcut 251-300 mg/dL - 7 units subcut 301-350 mg/dL - 10 units subcut 351-400 mg/dL - 12 units subcut  If blood glucose is greater than 400mg/dL, administer 14 units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat 10 units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. ONce blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale.
	<p><b>insulin aspart (Blank Insulin Aspart Sliding Scale)</b></p> <input type="checkbox"/> See Comments, subcut, inj, PRN glucose levels - see parameters If blood glucose is less than ___ mg/dL, initiate hypoglycemia guidelines and notify provider.  70-150 mg/dL - ___ units subcut 151-200 mg/dL - ___ units subcut 201-250 mg/dL - ___ units subcut 251-300 mg/dL - ___ units subcut 301-350 mg/dL - ___ units subcut 351-400 mg/dL - ___ units subcut  If blood glucose greater than 400 mg/dL, administer ___ units subcut, notify provider, and repeat POC blood sugar check in 90 minutes. Continue to repeat ___ units subcut and POC blood sugar checks every 90 minutes until blood glucose is less than 300 mg/dL. Once blood sugar is less than 300 mg/dL, repeat POC blood sugar in 4 hours and then resume normal POC blood sugar check and insulin aspart sliding scale.
<b>HYPOglycemia Guidelines</b>	
	<p><b>HYPOglycemia Guidelines</b></p> <input type="checkbox"/> ***See Reference Text***
	<p><b>glucose</b></p> <input type="checkbox"/> 15 g, PO, gel, as needed, PRN glucose levels - see parameters If 6 ounces of juice is not an option, may use glucose gel if blood glucose is less than 70 mg/dL and patient is symptomatic and able to swallow. See hypoglycemia Guidelines. Continued on next page....

TO  Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Physician Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_



SLIDING SCALE INSULIN ASPART PLAN

PHYSICIAN ORDERS

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER	ORDER DETAILS
	<p><b>glucose (D50)</b>  <input type="checkbox"/> 25 g, IVPush, syringe, as needed, PRN glucose levels - see parameters            Use if blood glucose is less than 70 mg/dL and patient is symptomatic and cannot swallow OR if patient has altered mental status AND has IV access. See hypoglycemia guidelines.</p>
	<p><b>glucagon</b>  <input type="checkbox"/> 1 mg, IM, inj, as needed, PRN glucose levels - see parameters            Use if blood glucose is less than 70 mg/dL and patient is symptomatic and cannot swallow OR if patient has altered mental status AND has NO IV access. See hypoglycemia guidelines.</p>

TO  Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Physician Signature: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

