UMC Health System				
NICU PROCEDURE PLAN		Patient I	Label Here	
	PHYSICIAN ORDERS			
Diagnosis				
Weight	Allergies			
	Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.			
ORDER	ORDER DETAILS			
	Patient Care			
	Confirm Central Line Placement (NICU) (Cleared for Use Central Line (NICU))			
	Line Type: PAL-Peripheral Arterial Line Line Type: UAC-Umbilical Artery Catheter	Line Type: PICC	us Catheter	
	Insert Central Line (NICU)			
	Line Type: PAL-Peripheral Arterial Line	Line Type: PICC		
		Line Type: UVC-Umbilical Veno	us Catheter	
	Insert Peripheral Line			
	Maintain Chest Tube			
	Protective Positioning (NICU)			
	Set Up for Lumbar Puncture			
	Set Up for Circumcision			
	Set Up for Chest Tube			
	Communication			
	Obtain Consent Consent for: PICC Line			
	Medications			
	Medication sentences are per dose. You will need to calculate a total daily dose if needed.			
	Iidocaine (lidocaine 1% preservative-free injectable solution)         0.5 mL, locally, inj, ONE TIME			
	petrolatum topical (petrolatum topical ointment) 1 app, topical, oint, 12x/day, PRN wound care			
	Apply with each diaper change.			
	acetaminophen (acetaminophen neonatal)			
	15 mg/kg, PO, liq, ONE TIME			
	***Do not exceed 2,600 mg of acetaminophen from all sources in 24 hours if under the age of 12 years. For all others do not			
	exceed 4,000 mg of acetaminophen from all sources in 24 hours***			
	Diagnostic Tests			
	DX Pedi Chest/Abd (DX Pedi Chest/Abd (NICU)) STAT, Line placement			
	DX Chest Portable			
	DX Abdomen Portable (DX Abdomen Portable (NICU))			
П то	Read Back	Scanned Powerchart	Scanned PharmScan	
Order Taken by Signature:		Date	Time	
Physician Signature:		Date	Time	