UMC Health System		Patient Label Here		
0	B/GYN MAGNESIUM SULFATE ADMINISTRATION P			
	PHYSICIAN ORDERS			
	Diagnosis			
Weight	Allergies			
	Place an "X" in the Orders column to designate orders of choice AN	VD an "x" in the specific order detail box(es) where applicable.		
ORDER	ORDER DETAILS Patient Care			
	Vital Signs ☐ Per Unit Standards, q15 min x 1 hour, then q30 min x 2 hours, then q	ı1h		
	Monitor Deep Tendon Reflexes			
	Patient Activity Bedrest	Bedrest   Up to Bedside Commode Only		
	Insert Urinary Catheter Foley, To: Dependent Drainage Bag			
	Urinary Catheter Care			
	Strict Intake and Output			
	Communication			
	Notify Provider of VS Parameters Temp Greater Than 100.4, RR Greater Than 28, RR Less Than 12, SpO2 Less Than 95, SBP Greater Than 160, SBP Less Than 90 Greater Than 110, DBP Less Than 50, HR Greater Than 120, HR Less Than 60			
	Notify Provider (Misc) ☐ Reason: Magnesium Level less than 4.8 mg/dL, or greater than 8.4 m respiratory depression)	ng/dL, or sign of toxicity (loss of deep tendon reflex, lethargy,		
	Notify Nurse (DO NOT USE FOR MEDS) Assess breath sounds every 2 hours			
	Seizure Precautions			
	Limit Stimuli			
	Limit Visitors			
	Dietary			
	NPO Diet			
	IV Solutions			
	***Total IV fluids are not to exceed 3 liters per 24 hours***			
	<b>LR</b> □ IV, 75 mL/hr	□ IV, 100 mL/hr		
	Medications			
	Medication sentences are per dose. You will need to calculate a tot ***A loading dose of 4-6 grams with a maintenance dose of 2-3 grams per impaired or weight of the patient is less than 45 kg***	-		
П то	Read Back	Scanned Powerchart Scanned PharmScan		
Order Take	en by Signature:	Date Time		
Physician Signature: Date Time				
		1201		

UMC Health System		Patient Label Here		
0	B/GYN MAGNESIUM SULFATE ADMINISTRATION PI	LAN		
	PHYSICIA	AN ORDERS		
	Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.			
ORDER	ORDER DETAILS			
	<ul> <li>magnesium sulfate</li> <li>4 g, IVPB, ivpb, ONE TIME, Infuse over 15 min, as loading dose.</li> <li>Repeat serum magnesium level 4 hours after the infusion is complete</li> <li>6 g, IVPB, ivpb, ONE TIME, Infuse over 15 min, as loading dose.</li> <li>Repeat serum magnesium level 4 hours after the infusion is complete</li> </ul>			
	magnesium sulfate 40 g/1,000 mL SW OB (magnesium sulfate 40 g/1         IV         magnesium sulfate 40 g/1,000 mL SW         Final Concentration: 40 mg/mL (0.04 g/mL)         FOR OB USE ONLY         Start at rate:         g/hr	1,000 mL SW - OB USE ONLY)		
	<b>calcium gluconate</b> 1 g, IVPB, ivpb, ONE TIME, PRN other, Infuse over 10 min For magnesium toxicity.			
	Laboratory			
	Magnesium Level ☐ Routine, T;N, q6h first draw 4 hours after magnesium loading dose. ☐ Routine, T;N, q12h first draw 4 hours after magnesium loading dose.			
	CBC with Differential	Routine, T;N, q12h		
	Comprehensive Metabolic Panel	Routine, T;N, q12h		
	Uric Acid Level			
	ALT Routine, T;N, q6h	Routine, T;N, q12h		
	AST Routine, T;N, q6h	Routine, T;N, q12h		
	LDH Routine, T;N, q6h	Routine, T;N, q12h		
	Prothrombin Time with INR			
	РТТ			
	Fibrinogen Level			
🗆 то	Read Back	Scanned Powerchart Scanned PharmScan		
Order Taken by Signature:		Date Time		
Physician Signature:				