

UMC Health System PEDIATRIC PICC INSERTION PLAN	Patient Label Here
----------------------------------------------------------------------	---------------------------

PHYSICIAN ORDERS

Diagnosis _____

Weight _____ **Allergies** _____

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER ORDER DETAILS

Patient Care

Intermittent Telemetry

Continuous Telemetry (Intermediate Care)

Continuous Pulse Oximetry

Consult RRT to Insert PICC Line

Communication

Notify Nurse (DO NOT USE FOR MEDS)
 Pre-Medicare patient as ordered prior to insertion.

Obtain Consent
 Consent for: PICC Insertion

Obtain Consent
 Consent for: Consent for: IV Sedation

Confirm Line Placement - Cleared for Use

IV Solutions

NS (NS bolus)
 20 mL/kg, IVPB, ONE TIME, Infuse over 30 min
 50 mL, IVPB, ONE TIME, Infuse over 1 hr
 200 mL, IVPB, ONE TIME, Infuse over 1 hr

20 mL/kg, IVPB, ONE TIME, Infuse over 60 min
 100 mL, IVPB, ONE TIME, Infuse over 1 hr

Medications

Medication sentences are per dose. You will need to calculate a total daily dose if needed.

lidocaine (lidocaine 1% preservative-free injectable solution)
 mL, transdermal, inj, ONE TIME

Pre Procedure Medication

acetaminophen (acetaminophen pediatric)
 15 mg/kg, per tube/PO, liq, ONE TIME, Give 30 minutes prior to procedure.
Give 30 minutes prior to procedure.

Do not exceed 2,600 mg of acetaminophen from all sources in 24 hours if under the age of 12 years. For all others do not exceed 4,000 mg of acetaminophen from all sources in 24 hour

LORazepam (LORazepam pediatric)
 0.02 mg/kg, per tube/PO, liq, ONE TIME, Give 30 minutes prior to procedure.
Give 30 minutes prior to procedure.
 0.03 mg/kg, per tube/PO, liq, ONE TIME, Give 30 minutes prior to procedure.
Give 30 minutes prior to procedure.
 0.05 mg/kg, IVPush, inj, ONE TIME, Give 30 minutes prior to procedure.
Give 30 minutes prior to procedure.
 0.1 mg/kg, IVPush, inj, ONE TIME, Give 30 minutes prior to procedure.
Give 30 minutes prior to procedure.
 0.1 mg/kg, IM, inj, ONE TIME, Give 30 minutes prior to procedure.
Give 30 minutes prior to procedure.

TO Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: _____ Date _____ Time _____

Physician Signature: _____ Date _____ Time _____



UMC Health System PEDIATRIC PICC INSERTION PLAN	Patient Label Here
----------------------------------------------------------------------	---------------------------

PHYSICIAN ORDERS

Place an "X" in the Orders column to designate orders of choice AND an "x" in the specific order detail box(es) where applicable.

ORDER	ORDER DETAILS
	midazolam (midazolam pediatric) <input type="checkbox"/> 0.5 mg/kg, PO, liq, ONE TIME, Give 10-30 minutes prior to procedure. Give 10-30 minutes prior to procedure. <input type="checkbox"/> 0.25 mg/kg, PO, liq, ONE TIME, Give 10-30 minutes prior to procedure. Give 10-30 minutes prior to procedure. <input type="checkbox"/> 0.1 mg/kg, IVPush, inj, ONE TIME, Give 10-30 minutes prior to procedure. Give 10-30 minutes prior to procedure.
	morphine (morphine pediatric) <input type="checkbox"/> 0.05 mg/kg, IVPush, inj, ONE TIME, Give 15-30 minutes prior to procedure. Give 15-30 minutes prior to procedure. Recommended maximum dose = 2 mg <input type="checkbox"/> 0.1 mg/kg, IVPush, inj, ONE TIME, Give 15-30 minutes prior to procedure. Give 15-30 minutes prior to procedure. Recommended maximum dose = 2 mg <input type="checkbox"/> 0.2 mg/kg, IVPush, inj, ONE TIME, Give 15-30 minutes prior to procedure. Give 15-30 minutes prior to procedure. Recommended maximum dose = 2 mg <input type="checkbox"/> 2 mg, IVPush, inj, ONE TIME, For patients weighing greater than or equal to 40 kg. Give 15-30 minutes prior to procedure. Give 15-30 minutes prior to procedure. <input type="checkbox"/> 4 mg, IVPush, inj, ONE TIME, For patients weighing greater than or equal to 40 kg. Give 15-30 minutes prior to procedure. Give 15-30 minutes prior to procedure.
PICC Line Flush	
	Use heparin flush 10 units/mL for patients weighing LESS than 18 kg heparin flush (heparin flush 10 units/mL injection (PEDI)) <input type="checkbox"/> 1 mL, IVPush, syringe, q12h, PRN flush, PICC, administer AFTER saline flush Flush PICC line with heparin solution every 12 hours even with a continuous infusion.
	Use heparin flush 100 units/mL for patients weighing GREATER than or EQUAL to 18 kg heparin flush (heparin flush 100 units/mL injection) <input type="checkbox"/> 1 mL, IVPush, syringe, q12h, PRN flush, PICC, administer AFTER saline flush Flush PICC line with heparin solution every 12 hours even with a continuous infusion.
Diagnostic Tests	
	DX Chest Single View <input type="checkbox"/> STAT, Portable, Line placement

TO Read Back

Scanned Powerchart

Scanned PharmScan

Order Taken by Signature: _____ Date _____ Time _____

Physician Signature: _____ Date _____ Time _____

