

Symptoms of HYPOglycemia

Mild:

- Diaphoretic
- Tachycardia
- Blurred Vision
- Sudden Weakness
- Shaking

Moderate:

- Dizziness
- Confusion
- Numbness of the lips
- Nausea/Vomiting
- Difficulty Swallowing

Severe: (May be combined with mild or moderate)

- Combative
- Seizures
- Loss of Consciousness

HYPOglycemia Guidelines

| SYMPTOMATIC PATIENT ABLE TO SWALLOW | | | |
|---|--|---|--|
| Blood Glucose Level | Treatment | Follow Up | |
| If blood glucose is less than 70 mg/dL and patient is symptomatic | Give 6 oz. of juice or 1 tube (15 gms) of glucose gel | Recheck blood glucose in 15 – 20 minutes Repeat treatment until blood glucose greater than 100 mg/dL Contact provider if blood glucose is less than 70 mg/dL for two consecutive measurements and patient is symptomatic. | |

| SYMPTOMATIC PATIENT UNABLE TO SWALLOW/NPO WITH IV ACCESS | | | |
|--|-----------------------------|--|--|
| Blood Glucose Level | Treatment | Follow Up | |
| If blood glucose is less than 70 mg/dL and patient is symptomatic and cannot swallow OR if patient has altered mental status | Dextrose 50% as ordered IV* | Recheck blood glucose in 15 – 20 minutes Repeat treatment until blood glucose greater than 100 mg/dL Contact provider if blood glucose is less than 70 mg/dL for two consecutive measurements and patient is symptomatic | |

SYMPTOMATIC PATIENT UNABLE TO SWALLOW/NPO WITHOUT IV ACCESS

| If blood glucose is less than | | |
|-------------------------------|--|--|
| 70 mg/dL and patient is | | |
| symptomatic and cannot | | |
| swallow OR if patient has | | |
| altered mental status | | |

Glucagon as ordered IM or Subcut*

- Contact physician for further orders
- Establish IV access
- Recheck blood glucose every 15 20 minutes

Use aspiration precautions as glucagon may cause nausea and vomiting

Source: UMC Pharmacy Version: 3 Reviewed: 12.20.2018

^{*} Dextrose and glucagon administration must be documented on the MAR